

**your
PR.i.VATES**

Chemotherapy

What is chemotherapy?

Surveillance vs one dose chemotherapy

What are the side affects of chemotherapy?

ORCHID 
FIGHTING MALE CANCER



www.yourprivates.org.uk



Contents

Introduction	3
Localised testicular cancer	4
Non localised testicular cancer	5
Cycles	6
Side effects	7

Chemotherapy

Chemotherapy is the use of anti-cancer (cytotoxic) drugs to destroy cancer cells. It works by disrupting the growth of cancer cells as they circulate in the bloodstream.

Localised Testicular Cancer

Early stage seminoma or non-seminoma which is confined to the testicle with no evidence of spread outside of it can usually be treated with a policy of surveillance.

This means regular CT scans and blood tests will be performed to detect any recurrence. Should testicular cancer reoccur then chemotherapy will be given and the chances are that a man will be cured.

However some men may find that they want to get on with their lives and not keep returning for scans as frequently as surveillance requires. In this situation a limited dose of chemotherapy can be given after surgery to reduce the risk of cancer returning.

If men opt for up front chemotherapy it will usually be a one off dose which may take around 1-2 hours to administer. Men may feel tired and suffer some minor side effects for a few weeks after.



Non Localised Testicular Cancer

For disease that has spread outside of the testicle a longer course of chemotherapy BEP (see below) will usually be suggested.

The drugs most commonly used to treat testicular cancer are Cisplatin, Etoposide and Bleomycin. A combination of all three is a treatment known as BEP. BEP chemotherapy can be given as a day case treatment or during a short stay in hospital. Blood tests will usually be performed prior to BEP to check that the body's immune system is healthy enough to cope with the treatment.


A Breathing test called a lung function test will be performed to ensure the lungs are functioning well as some types of chemotherapy may affect lung function.

A drip (cannula) will be sited into a vein in the hand, arm or neck and the chemotherapy administered through it. Anti-sickness (antiemetic), drugs can also be given through these tubes as well.

BEP chemotherapy can be given in different ways and the exact duration of treatment will be decided by an oncologist. It can be given as a regime over three days or five days on an inpatient or outpatient basis. The treatment regime is also known as a cycle. Each cycle of treatment is given over 3 weeks and 3 or 4 cycles of BEP chemotherapy are given depending on the extent of the disease.

Men who had high tumour markers or more widespread disease may be offered higher or different doses of chemotherapy.

Chemotherapy can temporarily cause infertility so men will be advised to store some sperm (sperm banking). Please see Testicular Cancer and Fertility Section for further information.



One cycle 3 Day BEP

(3-4 cycles over 2-3 months)

Day 1 Infusion of all BEP drugs

Day 2 Infusion of all BEP drugs

Day 3 Etoposide only

Day 8 Bleomycin

Day 15 Belomycin

Break

One cycle 5-day BEP

(3-4 cycles over 2-3 months)

Day 1 Etoposide and Cisplatin

Day 2 All BEP drugs


Day 3 Etoposide and Cisplatin

Day 4 Etoposide and Cisplatin

Day 5 Etoposide and Cisplatin

Day 15 Belomycin

Break



Side Effects

Below are some of the common and less common side effects associated with chemotherapy along with some tips that people have found useful on how to manage the symptoms.

Common side effects of chemotherapy:

Try:

Avoid:

Nausea and vomiting:

- Anti-sickness medication will be given on a regular basis and it is important to take this.
- Ginger, beer, tea or biscuits (eat regularly throughout the day).
- Peppermint tea, always drink slowly taking lots of sips.
- A small meal a few hours before chemo starts.


Never have chemotherapy on an empty stomach

- Hot and spicy foods (curry, Cajun cooking etc.).
 - Foods with high sugar content.
 - Fatty and greasy foods (chips, burgers etc.).
 - Large meals.
 - Eating and drinking too fast and drinking with meals.
 - Alcohol.
 - Caffeine (which is found in tea/ coffee/chocolate including chocolate bars/ energy drinks).
 - Baby food - this is designed for babies and has very little nutrition for adults!
-

Common side effects of chemotherapy:**Try:****Avoid:****Taste:**

(chemo can cause a metallic taste in the mouth)

- Fresh pineapple or other sharp tasting fruit.
- Boiled sweets while having treatment.
- Seasoned or marinated meat for meals to add flavour.
- Use herbs and spices in cooking to add stronger taste.

**Helpful tip**

Fresh pineapple or other sharp tasting fruit can help take away the metallic taste in your mouth.

See above for more tips

Mouth sores or ulcers from chemotherapy:

- Fresh pineapple can help prevent and heal mouth ulcers by stimulating saliva which protects the mouth.
- Bonjela.
- Soft child's/baby bristle toothbrush and baby/soothing (for instance) aloe vera toothpaste.
- Soft puréed or liquid diet to prevent chewing.
- Rinse the mouth with salt water on a regular basis if it can be tolerated; up to 4-5 times a day.
- Try sucking crushed ice during treatment.
- Artificial saliva which can be prescribed by a doctor.
- Vaseline for lips.
- Avoid tomato and citrus fruit based products and tobacco.
- Avoid oral care products that may cause a dry mouth and products that contain alcohol or peroxide.
- Hot and spicy foods (curry, Cajun etc.).
- Nuts and seeds.

**Common side effects
of chemotherapy:**
Try:
Avoid:
Tiredness

- Try and get small restful naps.
- Some people find benefit in Meditating.
- Sleeping tablets can be prescribed if insomnia becomes a problem.

Loss of appetite

- Try small frequent meals throughout the day.
- Try fortified soups/drinks or milk shakes.

Diarrhoea

- Rice and pasta.
- Dry crackers/biscuits.
- White bread.
- Bananas (a good source of potassium which is essential for the body's metabolism). Excessive diarrhoea will deplete
- potassium levels.

Hair Loss

(This may occur
2-3 weeks after
chemotherapy)

- Baseball cap.
- Shave hair off when it starts to fall out.
- Avoid tomato and citrus fruit based products and tobacco.
- Avoid oral care products that may cause a dry mouth and products that contain alcohol or peroxide.
- Hot and spicy foods (curry, Cajun etc.).
- Nuts and seeds.

**Low immune system
from chemotherapy**

- Pay meticulous attention to hygiene. Wash hands after going to the toilet!
- Travelling to foreign countries where risk of picking up illness is greater.
- Vaccines. Men will need to ask their oncologist if they will be able to travel and receive vaccines after treatment.

**Common side effects
of chemotherapy:**
Try:
Avoid:

**Tingling in the
arms and feet due
to nerve damage
(peripheral
neuropathy) from
chemotherapy**

- Keep hands and feet warm and avoid extremes of temperature.
- Take care when preparing food (use gloves) or when running hot water (test with an elbow) to prevent the risk of burning.
- Gentle exercise.

**Inflammation in the
lungs with shortness
of breath due to
low circulating
haemoglobin
(oxygenated blood)
from chemotherapy**

- A blood transfusion.
- Rest.
- Report any breathing difficulties.
- Avoid taxing, strenuous tasks.
- Do not smoke!

**Short of breath
with dizziness
due anaemia.
This is due to
low circulating
haemoglobin**

- If anaemia is severe may require blood transfusion.
- Eat green leafy vegetables and some red meat as these are rich in iron.
- Most nutrients can be obtained from eating normal food.
- Smoking, getting up too quickly as can cause sudden fainting.
- Avoid taking a lot of supplements without consulting a doctor.

**Tinnitus (ringing
in the ears) from
chemotherapy**

- Encourage people to speak slowly and clearly.
- Some people find that listening to gentle background music may help.

Common side effects of chemotherapy:**Try:****Avoid:****Skin changes**

- Electric shave to avoid cuts.
 - Non perfumed moisturizing Creams.
 - Chemo can make skin more sensitive to sunlight, check with the oncologist to see how long prolonged exposure to the sun should be avoided. Use factor 15 if sunbathing.
 - Allow 6 weeks after radiotherapy treatment before exposure to the sun it may be best to cover the treated area for up to a year.
 - Wear a hat in summer to avoid sun burn to the scalp.
- Wet shaving which may break the skin.
 - For radiotherapy it is useful to avoid non perfumed soaps and have tepid baths.
 - Avoid vigorous rubbing of the skin around the treated area.
 - Sun bathing in direct sun light as at risk of severe sun burn which can leave scarring for life.

**Helpful tip**

Some people find that non perfumed moisturising creams may help with skin changes.

See above for more tips



www.yourprivates.org.uk

A graphic for the Orchid National Male Cancer Helpline. It features a stylized white orchid flower icon on the left. To the right of the icon, the text "Orchid National Male Cancer Helpline" is written in a bold, white, sans-serif font. Below this, the phone number "0808 802 0010" is displayed in a large, bold, white, sans-serif font. At the bottom, the email address "helpline@orchid-cancer.org.uk" is written in a smaller, white, sans-serif font. The entire graphic is set against a dark blue background with a white border.